

Measuring the Proton-Proton Interaction Cross Section with Hybrid Data of the Pierre Auger Observatory

Olena Tkachenko
for Pierre Auger Collaboration


UHECR 2024
Malargue, Argentina



Cross sections for UHECR

$$\frac{dp}{dX_1} = \frac{1}{\lambda_{\text{int}}} e^{-X_1/\lambda_{\text{int}}}$$


$$\sigma^{\text{int}} = \frac{m_{\text{air}}}{\lambda_{\text{int}}}$$



Cross sections for UHECR

$$\frac{dp}{dX_1} = \frac{1}{\lambda_{\text{int}}} e^{-X_1/\lambda_{\text{int}}}$$


$$\sigma_{\text{int}} = \frac{m_{\text{air}}}{\lambda_{\text{int}}}$$






X_{max} distribution tail:

- $f(X_{\text{max}}) \sim e^{-X_{\text{max}}/\Lambda_\eta}$
- proton-dominated

Standard approach: proton-proton cross section from the tail fit




Standard approach: proton-proton cross section from the tail fit





*In original analysis (Phys. Rev. Let. 109, 2012) $\sigma_{\text{p-air}}$ was evaluated first and then converted into σ_{pp}

Standard approach: proton-proton cross section from the tail fit



He contribution \Rightarrow
systematic uncertainty



*In original analysis (Phys. Rev. Let. 109, 2012) $\sigma_{\text{p-air}}$ was evaluated first and then converted into σ_{pp}

Simultaneous Mass Composition and Cross Section Measurement

? Why?

Assumptions in the Standard Analyses:

- *Mass Composition:*
Assumes validity of a specific interaction model.
- *Interaction Cross Section:*
Relies on a proton-dominated tail of the X_{\max} distribution.

Simultaneous Mass Composition and Cross Section Measurement

? Why?

Assumptions in the Standard Analyses:

- *Mass Composition:*
Assumes validity of a specific interaction model.
- *Interaction Cross Section:*
Relies on a proton-dominated tail of the X_{\max} distribution.

💡 How?

- Vary the proton-proton cross section
- Perform a standard composition fit

Simultaneous Mass Composition and Cross Section Measurement

? Why?


Assumptions in the Standard Analyses:

- *Mass Composition:*
Assumes validity of a specific interaction model.
- *Interaction Cross Section:*
Relies on a proton-dominated tail of the X_{\max} distribution.

💡 How?

- Vary the proton-proton cross section
 - Perform a standard composition fit
- } → *self-consistent* estimation of the interaction cross sections and cosmic-ray primary composition

Rescaling of the interaction cross section



Rescale the proton-proton cross section:

$$\sigma_{\text{mod}}^{\text{PP}} = \sigma_{\text{orig}}^{\text{PP}} f^{\text{PP}}(E_0, E),$$


with a linear scaling factor $f^{\text{PP}}(E_0, E)$:

$$f^{\text{PP}}(E_0, E) = 1 + H(E - E_0) (f_{\text{lg } E_1}^{\text{PP}} - 1) \frac{\lg(E/E_0)}{\lg(E_1/E_0)}.$$

Same approach for cross section modification as in original analysis^{1,2} but instead of $\sigma_{\text{p-air}}$ we directly modify σ_{pp}

* H is a Heaviside function;

* $f_{\text{lg } E_1}$ is the rescaling factor at $E = E_1$ (here $E_1 = 10^{19}$ eV);



* Threshold E_0 sets an energy above which cross sections are modified;


* Modified implementation of Sibyll 2.3d hadronic interaction model.

¹ Phys. Rev. Lett. 109 (2012) 062002.

² R. Ulrich et al, Phys. Rev. D 83 (2011) 054026 .


Algorithm: fitting procedure

- 1 get the X_{\max} distributions for the discrete set of $f_{\lg E}^{\text{pp}}$ values;




Algorithm: fitting procedure

- 1 get the X_{\max} distributions for the discrete set of $f_{\lg E}^{\text{pp}}$ values;
- 2 perform the 4-component binned maximum likelihood mass composition with for varied:
 - rescaling factor $f_{\lg E}^{\text{pp}}$ [0.2, 3.0]
 - shift in the X_{\max} [-50, 40] g/cm²




Algorithm: fitting procedure

- 1 get the X_{\max} distributions for the discrete set of $f_{\lg E}^{\text{pp}}$ values;
- 2 perform the 4-component binned maximum likelihood mass composition with for varied:
 - rescaling factor $f_{\lg E}^{\text{pp}}$ [0.2, 3.0]
 - shift in the X_{\max} [-50, 40] g/cm²




Algorithm: fitting procedure

- 1 get the X_{\max} distributions for the discrete set of $f_{\lg E}^{\text{pp}}$ values;
- 2 perform the 4-component binned maximum likelihood mass composition with for varied:
 - rescaling factor $f_{\lg E}^{\text{pp}}$ [0.2, 3.0]
 - shift in the X_{\max} [-50, 40] g/cm^2
- 3 sum χ^2 for each δX_{\max} and $f_{\lg E}^{\text{pp}}$ over the considered energy range;



Algorithm: fitting procedure

- 1 get the X_{\max} distributions for the discrete set of $f_{\lg E}^{\text{pp}}$ values;
- 2 perform the 4-component binned maximum likelihood mass composition with for varied:
 - rescaling factor $f_{\lg E}^{\text{pp}}$ [0.2, 3.0]
 - shift in the X_{\max} [-50, 40] g/cm^2
- 3 sum χ^2 for each δX_{\max} and $f_{\lg E}^{\text{pp}}$ over the considered energy range;
- 4 find the best-fit χ^2 and get the corresponding cross section σ_{pp} , shift δX_{\max} , and composition.



Systematic uncertainties

Origin	Impact on σ_{pp} , %	Impact on δX_{\max} , g/cm ²
Energy scale	-3.1	+6 -4
Detector effects	+7 -12	± 1
E -dependent X_{\max} syst.	± 2	± 7
Composition	+3 -7	+5
Elasticity	+15 -17	+1 -3
Multiplicity	+9	+1 -8



Under evaluation: mass-dependent shift in X_{\max}

Decrease in uncertainty compared to standard analyses:



- up to 25% He-fraction bias;
- X_{\max} scale systematics.

Composition-related bias



Simulations for $10^{17.8}$ - $10^{17.9}$ eV energy bin



Composition-related bias





- Lowest E : $N \sim 10^4$
- Highest E : $N \sim 50$
- Considering the χ^2 sum over all energies compensates for the possible bias contribution from the highest energies



*fits are shown for one energy bin

Why fitting a shift in X_{\max} matter


Fit with the modified Sibyll 2.3d to QGSJETII.4 and Sibyll 2.1 simulations




*The fitted X_{\max} scales are close to the differences between the models

Why fitting a shift in X_{\max} matter

Fit with the modified Sibyll 2.3d to QGSJETII.4 and Sibyll 2.1 simulations




Allowing for the shift in the X_{\max} scale to vary in the fit enables accurate recovery of the cross section and composition when the scale differs from the Sibyll 2.3d



*The fitted X_{\max} scales are close to the differences between the models

Analysis Results



Simulations with AugerMix



*AugerMix==composition as observed in Auger data

Analysis Results

Simulations with AugerMix





*AugerMix==composition as observed in Auger data

*statistical uncertainty on the mass & xsec fit is from averaging the 100 sim. data realizations

Analysis Results

Simulations with AugerMix, including possible systematics



*AugerMix==composition as observed in Auger data
 *statistical/systematical uncertainties correspond to one sim. data realization



Summary

Simultaneous estimation of the cosmic-ray mass composition and proton-proton interaction cross section:

- Remove degeneracy from previous analyses:
 - independent on the underlying composition;
 - independent on the underlying cross section;
- Improvement in the statistical/systematic uncertainty compared to standard analyses:
 - suppressed He fraction-related systematics
 - suppressed X_{\max} scale systematics
- Higher confidence in the estimation;


Future plans

- Careful evaluation of the potential biases and systematics
- Apply to the full Phase I Auger data



Back-up



Mass composition from data



*default cross section and X_{max} scale

Energy-dependent X_{\max} systematics


Distribution of fit results assuming the energy-dependent shifts in X_{\max} scale



*100 simulated data realizations



Energy-dependent X_{\max} systematics

Distribution of fit results assuming the energy-dependent shifts in X_{\max} scale





*100 simulated data realizations

Mass composition measurements: effect of cross section rescaling



Rescaling by $\pm 20\%$ at 10^{19} eV

Fit results for the different composition scenarios



*Constant composition over the considered energy range
 *Number of events in each energy bin the same as in Auger data